



Catry, P., Campos, A. R., Granadeiro, J. P., Neto, J. M., Ramos, J., Newton, J., and Bearhop, S. (2016) Provenance does matter: links between winter trophic segregation and the migratory origins of European robins. *Oecologia*, 182(4), pp. 985-994. (doi:10.1007/s00442-016-3725-z)

This is the author's final accepted version.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/129964/>

Deposited on: 07 November 2016

PROVENANCE DOES MATTER: LINKS BETWEEN WINTER TROPHIC

SEGREGATION AND THE MIGRATORY ORIGINS OF EUROPEAN ROBINS

Paulo Catry¹, Ana R. Campos², José Pedro Granadeiro³, Júlio M. Neto^{4,5}, Jaime Ramos², Jason Newton^{6,7}, Stuart Bearhop⁷

¹ MARE – Marine and Environmental Sciences Center, ISPA - Instituto Universitário, Rua Jardim do Tabaco 34, 1149-041 Lisboa, Portugal. Email: paulo.catry@gmail.com

² MARE – Marine and Environmental Sciences Centre, Department of Life Sciences, University of Coimbra, 3004-517 Coimbra, Portugal.

³ CESAM and Departamento de Biologia Animal, Faculdade de Ciências da Universidade de Lisboa, Lisboa, Portugal.

⁴ CIBIO/UP — Centro de Investigação em Biodiversidade e Recursos Genéticos, Universidade do Porto, Portugal

⁵ Molecular Ecology and Evolution Lab, MEMEG, Department of Biology, University of Lund, Sweden

⁶ NERC Life Sciences Mass Spectrometry Facility, SUERC, Rankine Avenue, East Kilbride, G75 0QF, UK.

⁷ Centre for Ecology and Conservation, University of Exeter, Cornwall Campus, TR10 9FE, UK.

PC, ARC conceived and designed the experiments, PC, ARC, JMN, JN performed the experiments, PC, ARC, JPG analysed the data, PC, ARC, JR, SB wrote the manuscript.

24 *Abstract.*

25 Amongst migratory species, it is common to find individuals from different populations or
26 geographical origins sharing staging or wintering areas. Given their differing life histories,
27 ecological theory would predict that the different groups of individuals should exhibit some level
28 of niche segregation. This has rarely been investigated because of the difficulty in assigning
29 migrating individuals to breeding areas. Here, we start by documenting a broad geographical
30 gradient of hydrogen isotopes ($\delta^2\text{H}$) in robin *Erithacus rubecula* feathers across Europe. We then
31 use $\delta^2\text{H}$, as well as wing-tip shape, as surrogates for broad migratory origin of birds wintering in
32 Iberia, to investigate the ecological segregation of populations. Wintering robins of different
33 sexes, ages and body sizes are known to segregate between habitats in Iberia. This has been
34 attributed to the despotic exclusion of inferior competitors from the best patches by dominant
35 individuals. We find no segregation between habitats in relation to $\delta^2\text{H}$ in feathers, or to wing-tip
36 shape, which suggests that no major asymmetries in competitive ability exist between migrant
37 robins of different origins. Trophic level (inferred from nitrogen isotopes in blood) correlated
38 both with $\delta^2\text{H}$ in feathers and with wing-tip shape, showing that individuals from different
39 geographic origins display a degree of ecological segregation in shared winter quarters. Isotopic
40 mixing models indicate that wintering birds originating from more northerly populations
41 consume more invertebrates. Our multi-scale study suggests that trophic-niche segregation may
42 result from specializations (arising in the population-specific breeding areas) that are transported
43 by the migrants into the shared wintering grounds.

44
45 Keywords: Seasonal matching, ecological segregation, *Erithacus rubecula*, migration, stable
46 isotopes

INTRODUCTION

Billions of migratory birds, as well as countless other animals, seasonally shift positions across vast geographical areas. At the end of the migratory journey, they often face numerous conspecific competitors from diverse geographical origins that share common non-breeding quarters (Block et al 2005, Newton 2008, Blumenthal et al 2009). The degree of population overlap and competition in the non-breeding season may have varied implications, from demographic regulation to the evolution and shaping of migratory systems (Rickleffs 1992, Bell 2005), but because of difficulties in assigning individuals to different origins, thus far virtually nothing is known of the ways individuals from different populations of the same species coexist.

There are theoretical reasons to expect segregation and ecological differentiation when multiple populations locally coexist. For example, local adaptations or specializations associated with the breeding grounds in different latitudes and habitats (Peterson 1993) may well be transported to wintering sites by migrating individuals (Price & Gross 2005, Rayner et al. 2011). Alternatively, competitive asymmetries on the wintering grounds may force habitat/trophic segregation of different populations where resources are limited (Fretwell & Lucas 1970, Greenberg 1986). For example, if some migrants arrive earlier, they may gain advantage from earlier ownership of non-breeding territories and win more territorial contests against later arrivals (Tobias 1997). In this scenario, populations at disadvantage may have no alternative but occupying different niches (Greenberg et al. 2001), perhaps with time adapting to them. Asymmetries may also occur due to carry-over effects (Catry et al. 2013) and matching of habitat quality between breeding and wintering sites has also been reported for non-territorial migrants (Gunnarsson et al. 2005).

In species such as European robins *Erithacus rubecula* (hereafter “robins”) and in blackcaps *Sylvia atricapilla*, sedentary Iberian populations mostly occupy woodland habitats where they also nest, whereas migrants predominate at shrubland sites where no breeding takes place (Pérez-Tris & Tellería 2002, Tellería & Pérez-Tris 2004). It is unclear whether such segregation results from the predominance of sedentary individuals at sites where they hold year-round territories (Campos et al. 2011a), or whether there is a difference in winter habitat choice irrespective of its potential value for future reproduction. A recent study suggests that amongst migratory blackcaps there is no difference in the habitat distribution of birds of different origins (de la Hera et al. 2012). However, segregation may also occur at finer ecological scales, such as among micro-habitats or dietary preferences.

In this paper, we use robins as a model species to determine whether there is an ecological segregation of wintering migratory individuals of different geographical origins at multiple ecological scales. We used stable hydrogen isotope ratios ($\delta^2\text{H}$) in robin feathers to give an indication of geographic origin, as these ratios in amount-weighted mean annual precipitation occur in a pronounced gradient across Europe along a southwest-northeast axis (Hobson et al. 2004). This is the main direction of migration of robins occurring in western Iberia (Bueno 1998). We also use stable carbon and nitrogen isotope ratios and multiple source mixing models to assess differences in diet (Inger & Bearhop 2007, Parnell *et al* 2010). Specifically we investigate the question at three hierarchical spatial scales plus at one further ecological dimension: (1) regional, by comparing different sectors of Iberia; (2) macro-habitat, by comparing two contrasting habitats in the same region; (3) micro-habitat, by making detailed habitat measurement within two study sites; and (4) diet, as assessed through stable-nitrogen isotope ratios in blood, a proxy for trophic level.

92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114

METHODS

Study species

European robins are small passerines common across almost the entire continent. Northern and eastern populations are fully migratory, while at the western and southern edge of the range (including much of Iberia) they are mostly sedentary. At mid-latitudes, robins are partial migrants (Cramp 1988). In western Iberia, robins are extremely common winter visitors, occupying virtually every type of habitat with some tree or shrub cover, feeding on invertebrates, berries, olives and acorns (Herrera 1977, 1998). Some individuals defend winter territories while others behave as floaters (Cuadrado 1997).

Feather sampling across Europe

Because feathers, once fully developed, are metabolically inert, they reflect the environment in which they were grown. Flight feathers of robins <1 year old are grown in the nest. Adult feathers are moulted after reproduction, while birds are still on the breeding grounds (Cramp 1988). Note that robins are highly faithful to nesting areas and breeding dispersal, when it occurs, is mostly of less than 10km (Paradis et al. 1998). We sampled robin feathers grown in 2007 across multiple European locations in Portugal, France, Sweden and Russia. Stable isotope ratios in terrestrial ecosystems are known to show site-specific variations that are superimposed on broad geographical trends (Wunder et al. 2005). Hence, to avoid obtaining spurious results caused by local variation, within each region/country we tried, as much as possible, to sample individuals from as many different locations (covering a broad range of latitudes) as possible. Adult robins were sampled during the 2008 breeding season in Portugal (16 sites from north to

south; late April to mid July), France (10 sites, from north to centre, late May to early July) and Russia (2 sites, Kaliningrad enclave, Baltic region, late May and June). Scandinavian robins were sampled during migration at the southern tip of Sweden, at Falsterbo (55° 23' N, 12° 50' E), where migrant robins originating from all over Sweden and (less so) from Finland occur during autumn (Karlsson et al. 1988). Trapping took place on 5 different dates, from mid September to mid-October 2007, covering the main migratory period. It is therefore likely that each of the 36 individuals sampled here came from a different breeding site. Note that individuals sampled in autumn 2007 and adults sampled in spring 2008 would have all grown their feathers in the same breeding season.

Each sampled robin was ringed. The innermost secondary was cut and preserved for isotope analysis. This feather was chosen as it is rarely lost by accident, which prevents the unintended sampling of replacement feathers grown away from the breeding site.

Winter sampling in Iberia

Robins were captured using mist-nets and baited traps during the winter (from mid-November 2008 to mid-February 2009) at 5 locations, covering the full wintering range along a NE-SW axis on the western part of Iberia: (1) Irun, Northern Spain (43°20'N, 1° 47'W) in mostly woodland habitats (10m altitude; N = 22 robins); (2) Salvaterra do Extremo, Central Portugal (39° 52'N, 06° 54' W) in open cork oak woodland (350m altitude; N = 12 robins); (3) Charneca (near Alcochete), Central/South Portugal (38° 49'N, 08° 49' W) in open cork oak woodland with virtually no undergrowth (25m altitude; N = 45 robins); (4) Arrábida, Central/South Portugal (38° 27'N, 09° 01' W) in Mediterranean shrubland (200m altitude; N = 77 robins); (5) Charito, Algarve, South Portugal (37°10'N, 08° 27'W), in an open mixture of woodland/shrubland with

carob trees, mastic and olive trees (50m altitude; N = 17 robins). Robins do not nest at any of the sites where traps or nets were placed.

Sites 3 and 4 are separated by just 45km and have a similar climate. They were selected to represent the greatest possible contrast in terms of shrub and tree cover, as these variables are known to influence the distribution of robin of different age, sex and size classes (Tellería *et al.* 2001, Catry *et al.* 2004, own unpubl. data). Within these main study sites (sites 3 & 4), robins were trapped with spring traps baited with mealworms only, to ensure they were foraging at the site of capture, and not just commuting to another place. At each exact capture location (and following Catry *et al.* 2004) we visually estimated the following micro-habitat variables, in a radius of 20m around the trap: a) mean canopy height, b) percentage canopy cover, c) mean height of the shrub layer (shrubs were defined as woody plants less than 3 m tall), d) percentage shrub cover, e) percentage bare ground or short-grass (< 10 cm tall) cover and f) percentage long-grass cover.

Robins captured in winter were aged according to Svensson (1992) and processed in the same way as described above for spring. Additionally, for birds captured at Charneca and Arrábida, we measured each of the 9 primaries using a ruler with a pin. Pectoral muscle profiles were scored with reference to the prominence of the sternal keel and muscle shape, on a 4-point scale following Bairlein (1995; see also Gosler, 1991).

Approximately 60 µl of blood were obtained by puncturing the *vena ulnaris*. A drop was preserved in ethanol for molecular sexing and the remaining was frozen until further processing. Sexing was done through the amplification by PCR (polymerase chain reaction) of a fragment of the CHD gene, using the primers P2 and P8 (Griffiths *et al.* 1998).

At Charneca and Arrábida, samples of the most common invertebrates, berries and acorns likely to be part of robin diet (Herrera 1977, 1998, Debusse & Isenmann 1985) were collected during winter and frozen for subsequent laboratory analysis.

Stable isotopes

i) Hydrogen isotopes

The sampled feathers from our study birds were carefully washed in distilled water and oven dried and left for a week equilibrating. Feathers were then cut very finely and weighed into silver cups and loaded into a zero-blank autosampler ready for analysis by continuous flow isotope ratio mass spectrometry (CF-IRMS), using a high temperature reduction system (TC/EA) interfaced with a Thermo-Fisher-Scientific Delta V Plus IRMS. We used the “comparative equilibration” (Wassenaar and Hobson, 2003) approach to correct the $\delta^2\text{H}$ isotope data, Briefly, the steam equilibration takes place in a modified Costech zero-blank autosampler containing keratin standards. Water of known $\delta^2\text{H}_{\text{VSMOW}}$ (W64444, $\text{d}2\text{H}=-399.1\text{‰}$, provided by USGS-RSIL; or alternatively S9, an internal water standard, $\delta^2\text{H}_{\text{VSMOW}} = +99.09\text{‰}$) is injected into the evacuated hot autosampler in a 110°C oven. Following equilibration, evacuation and cooling to room temperature the autosampler is reattached to the TC/EA, purged with helium gas and run as normal. Non-exchangeable standards are used to correct raw hydrogen isotope data; these were IAEA-CH7 (polyethylene foil, $\delta^2\text{H}_{\text{VSMOW}} = -100.3\text{‰}$), C36 (hexatriacontane provided by A. Schimmelmann, $\delta^2\text{H}_{\text{VSMOW}} = 246.7\text{‰}$) and an internal polythene foil standard “BOS” ($\delta^2\text{H}_{\text{VSMOW}} = -82.1\text{‰}$). Two runs using the two widely disparate water standards above, allows %H exchangeability and unexchangeable $\delta^2\text{H}_{\text{VSMOW}}$ to be calculated. As such that the $\delta^2\text{H}$ data here represent the unexchangeable hydrogen and fixed to the VSMOW-VSLAP scale.

The standards used for this approach are CFS (chicken feathers; $\delta^2\text{H}_{\text{VSMOW}} = -147 \pm 5 \text{ ‰}$, Hobson and Wassenaar 2008), BWB-II (bowhead whale baleen; $\delta^2\text{H}_{\text{VSMOW}} = -108 \pm 4 \text{ ‰}$, Hobson and Wassenaar 2008) and ISB (Icelandic black-legged kittiwake, *Rissa tridactyla* feathers $\delta^2\text{H}_{\text{VSMOW}} = -72 \pm 3 \text{ ‰}$, Fox et al. 2007). A fourth keratin standard WG (Willow grouse *Lagopus lagopus* feathers, $\delta^2\text{H}_{\text{VSMOW}} = -135 \pm 1 \text{ ‰}$, unpublished data) is run independently of the calibration (see also Evans et al. 2012). All four standards were run in triplicate in five measurement runs; collectively the fifteen WG standards had a $\delta^2\text{H}_{\text{VSMOW}}$ of $-135.38 \pm 2.89 \text{ ‰}$. All $\delta^2\text{H}$ values quoted are non-exchangeable hydrogen derived from the comparative steam equilibration technique.

ii) Carbon and nitrogen isotopes

Whole blood, whole insects, berries and acorns were oven dried, reduced to a homogeneous powder, and weighed into tin cups then also analysed for nitrogen ($\delta^{15}\text{N}_{\text{AIR}}$) and carbon ($\delta^{13}\text{C}_{\text{V-PDB}}$) isotope ratios by CF-IRMS, using a Costech ECS 4010 elemental analyser connected to a Thermo-Fisher-Scientific Delta XP Plus IRMS. Internal laboratory standards included gelatin, and two alanine standards spiked with ^{13}C - and ^{15}N -enriched alanine respectively. These internal standards are checked monthly against international reference materials USGS 40 and USGS 41 (glutamic acid), IAEA N1, N2 and USGS25 (ammonium sulphate) and IAEA CH6 (sucrose). Repeated (>20) $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ measurements in each run gave standard deviations of ~ 0.2 and 0.1 ‰ respectively.

All the analyses were performed at the same laboratory of the NERC Life Sciences Mass Spectrometry Facility at East Kilbride, Scotland.

Statistical analysis

An index of shrub volume at each trapping location was obtained by multiplying shrub cover by its mean height. This micro-habitat variable has been shown to contribute to predict size, age and sex-ratios of captured robins in winter (Catry et al. 2004).

Bird populations with a greater migratory tendency tend to have longer and more pointed wings, and this general pattern applies to robins, with more northerly populations differing from southern and predominantly sedentary ones (Pérez-Tris *et al.* 2000, Tellería *et al.* 2001). Hence, using the measurements of individual primary feathers, we calculated a wingtip pointedness index (C_2) derived from Size-Corrected Components Analysis - SCCA (Lockwood *et al.* 1998). This was only done for robins wintering at Charneca and Arrábida, as most other samples had been collected by collaborators who were unable to provide measurements of primary feathers.

In order to estimate the contribution of each prey type in the diet of individual robins, we developed an isotopic mixing model in the “R” package SIAR (Parnell et al. 2010, R Development Core Team 2012). The procedure implements a Markov Chain Monte Carlo algorithm to seek likely combination of prey contributions that best match the isotopic signature of the consumer, taking into account tissue- and species-specific discrimination factors of dietary items. Natural variability in isotopic signature of food sources and in trophic discrimination is dealt with within the Bayesian approach implemented by SIAR (Parnell *et al.* 2010). Prey items were aggregated according to prior knowledge of their feeding habits in the regions and included four major types: Ants, other insects, *Quercus sp.* acorns and berries, the latter only present in one of the sites (Arrábida). Trophic enrichment factors were set at $1.7 \pm 0.17\text{‰}$ and $2.4 \pm 0.24\text{‰}$ and for carbon and nitrogen respectively (Hobson and Bairlein 2003).

We used general linear models to investigate the effects of single factors/variables on stable hydrogen isotope ratios or C_2 , and checked their significance using likelihood ratio tests

against the corresponding null models (i.e. with the variable of interest removed). We also examined the extent to which $\delta^{15}\text{N}$ in blood was influenced by age, sex, month, habitats and either $\delta^2\text{H}$ or C_2 (as indicators of origin), for which general linear models within a Bayesian model averaging (BMA) framework were used. Variables were checked for normality and homogeneity of variance. To avoid over-fitting we tested all combinations of variables, but only included first-order interactions between month, age, sex and study site. Computations were carried out using function `bic.glm` of BMA package (Raftery et al. 2013), assuming uniform priors for the model parameters and using the Bayesian Information Criterion (BIC) approximation to estimate the integrated model likelihood. We only averaged a subset of all possible models, by constraining the ratio of the posterior probabilities of the candidate model to the best model to be higher than 0.05 (Occam's window criteria Madigan and Raftery 1994). For models including either $\delta^2\text{H}$ or C_2 we calculated the posterior probability that each variable is in the model (in %), as well as the posterior mean and standard deviation for all estimated coefficients.

RESULTS

Hydrogen isotope trends in Europe

There were highly significant differences in $\delta^2\text{H}$ values in feathers sampled across a range of different European countries (GLM log-likelihood ratio $\chi^2 = 16499$, $P < 0.001$), with a clear trend for more negative values with increasing distance from Portugal towards north-eastern Europe (Fig. 1). The patterns for the more extensive Portuguese data set are similar, illustrating that even within a smaller region, closer to our winter study sites, latitude of a site and mean $\delta^2\text{H}$

values of robins from that same location correlate significantly ($r = -0.67$, $N = 16$ sites, $P = 0.005$).

Segregation in winter quarters

As expected there was a correlation between wing pointedness (C_2) and δ^2H in feathers ($r = 0.38$, $N = 121$, $P < 0.001$) for birds captured in central Portugal during winter, indicating that birds with more pointed wings had bred (or were hatched) further northeast. Using the larger data set of birds captured across Iberia, wing length (no data for C_2) also correlated with δ^2H ($r = -0.27$, $N = 170$, $P < 0.001$), again indicating that birds with longer wings originated further northeast.

The δ^2H in feathers did not correlate with latitude of the Iberian wintering site ($r = 0.03$, $N = 174$, $P = 0.69$). Although this analysis is pseudoreplicated (as many individuals came from the same wintering site), we choose to present it as a conservative example of there being no relationship between breeding latitude and wintering latitude. Furthermore, as shown below, there are no reasons to expect important site-related differences in δ^2H composition of feathers of *wintering* robins. The correlation between mean δ^2H in robin feathers from a given wintering site and latitude is also not significant ($r = 0.33$, $N = 5$, $P = 0.59$).

The feathers of robins wintering in open woodland and in shrubland in nearly adjacent sites in central Portugal did not differ significantly in δ^2H (respectively, $-87.4 \pm 9.6\text{‰}$, $N = 45$ and $-86.1 \pm 12.6\text{‰}$, $N = 77$, GLM log-likelihood test $\chi^2 = 0.002$, $P=0.997$). None of the microhabitat variables measured at Arrábida and at Charneca correlated with stable hydrogen isotope ratios or with the wing pointedness index (all correlations with $P > 0.05$, $N = 45$ at Charneca and $N = 77$ at Arrábida).

Model averaging revealed that values of $\delta^{15}\text{N}$ in whole blood differed between habitats, among sampling months and were negatively correlated with $\delta^2\text{H}$ (Table 1, Fig. 2), while age and sex effects and interactions were non-significant. The same highly significant relationship holds when $\delta^2\text{H}$ replaces the wing pointedness index C_2 (Table 1, Fig. 2). The percentage contribution of invertebrates in the diet of individual birds (estimated with isotopic mixing models; see Electronic Supplementary Materials for data on Carbon and Nitrogen isotope ratios in prey and robins) showed a negative relationship with $\delta^2\text{H}$ levels (GLM, $t = -2.16$, $P = 0.034$), and also a significant effect of site (GLM, $t = 14.6$, $P < 0.001$).

DISCUSSION

For the first time we have systematically measured variation across multiple niche dimensions amongst sympatrically wintering conspecific migrants from allopatric breeding areas. Perhaps surprisingly we found that robins from a wide range of breeding latitudes within Europe show no evidence of geographic structuring or migratory connectivity within Iberia. Likewise we could find no consistent differences in habitat and microhabitat selection. However, individuals from different origins differed in their trophic position during the winter months, suggesting a form of seasonal matching (Gunnarsson et al. 2005) between the region of birth or reproduction and the wintering ecology of migrants. Trophic specialisation of this nature may play a key role in allowing these populations with different migratory tendencies to co-exist during the non-breeding season.

We began by documenting a clear geographical cline of hydrogen isotope ratios in robin feathers across Europe. The relationship between wing length and shape and $\delta^2\text{H}$ in wintering birds is also in line with the initial expectation of more negative ratios for birds with longer and

more pointed wings, which likely belong to more migratory and northerly distributed populations (Pérez-Tris et al. 2000).

The distribution of $\delta^2\text{H}$ values evidenced in Fig. 1 clearly indicates that the robin communities wintering in our study sites are a mixture of individuals originating from a wide range of latitudes in Europe (a condition necessary for our study) and is in agreement with previous research (Bueno 1998, Catry et al. 2010, Campos et al. 2011a). This panmitic overwintering model at the level of the Iberian Peninsula is similar to that found on overwintering monarch butterflies (Wassenar & Hobson 1998).

Many conspecific migrant populations show some segregation in winter quarters, a phenomenon known as migratory connectivity (Newton 2008). Segregation by habitat has more rarely been documented (Marion 1995, Duijns et al. 2009, Alves et al. 2010). In robins, however, we found no habitat influence on the geographical composition of wintering birds, despite the fact that we compared contrasting habitats which, for example, harbour robin communities with markedly different sex-ratios (Catry et al. 2004 and own unpubl. data). A similar result has been obtained for blackcaps, a mostly non-territorial passerine wintering in southern Iberia (de la Hera et al. 2012).

Amongst migrants that are territorial during winter, asymmetries in resource-holding potential may result in a despotic distribution, where dominant individuals relegate weaker competitors to less suitable habitats (e.g. Marra 2000). Detailed studies of wintering robins in Iberia have indicated that local birds may have a competitive advantage over visiting migrants, given that they occupy the putative best habitats where they nest throughout the year (Tellería et al. 2001, Tellería & Pérez-Tris 2004). However, the same patterns of population segregation might not necessarily arise from the competitive superiority of southern individuals, but rather

from them opting to invest in territorial defence over winter at appropriate sites, in preparation for future breeding attempts. The present study was conducted only at sites where no robins nest. Under these conditions, there was no evidence for habitat separation between birds from different origins, which would be expected if robins from different latitudes had dissimilar abilities to compete for the best wintering sites. Experimental work with artificial feeders also suggested no differences in the resource holding potential linked to geographical origin of robins (Campos et al. 2011b).

When potentially competing groups do not segregate in one niche dimension, they are more likely to differ in another one, and this is exactly what we found, with robins from more northerly origins feeding at a higher trophic level (consuming more invertebrates). Wintering robins are broadly omnivorous (Herrera 1977, 1978, 1998, Debusse & Isenmann 1985), which means that any dietary specialization or differentiation is likely to be picked up by an analysis of trophic level given by an indicator such as $\delta^{15}\text{N}$ in blood. Most robins arrive in October (Bueno 1998, Catry et al. 2010), and given the fact that turnover half-life of whole blood for birds the size of a robin is ca. 5 days (Hobson 2008), it is almost impossible that any isotopic measurements made during late-November to February retain any signature of the migratory habitats or of the migratory journey itself.

Virtually no studies have documented dietary segregation of coexisting migratory populations in common wintering grounds, and the few exceptions reported very small differences between distinct subspecies (Duijns et al. 2009, Fonteneau et al. 2009). An exception is a study of black-tailed godwits *Limosa limosa* that documented important dietary differences between subspecies, but those were linked to a clear segregation by habitat (Alves et al. 2010). A study of collared flycatchers *Ficedula albicollis* using isotopes also suggested there may be

consistent interpopulation differentiation in the feeding ecology at the wintering sites, but alternative explanations for the patterns found could not be rejected (Hjernquist et al. 2009). Individual specialization in diet has been suggested for robins of multiple populations wintering in southern Iberia, which was attributed to differences in morphology, but those were not linked to geographical origin of migrants (Herrera 1978). Our study provides the first strong evidence that different migratory populations may have different diets, even when sharing wintering habitats.

Robins from different breeding origins may have slightly divergent adaptations, allowing them to better exploit the resources of their nesting habitats, resulting in an intraspecific niche differentiation (Peterson & Holt 2003). Feeding specializations have been documented in passerine birds even at small spatial scales (10s of kilometres, along an altitudinal gradient) and despite gene flow (e.g. McCormack & Smith 2008). Hence, it seems plausible to speculate that robins originating from different parts of the wide European range could carry different morphological, physiological or behavioural specializations, resulting from local adaptations, into the winter quarters (see Price & Gross 2005, Rayner et al. 2011 for this type of phenomenon amongst closely related migratory species and subspecies). This would then lead them to adopt different diets when sharing a common environment. Such specializations could even be reinforced by selection during winter-time, if they were to reduce competition for resources. An alternative explanation would be that longer-distance migrants have different nutritional requirements than shorter-distance migrant conspecifics (Tellería et al. 2013), but with the present knowledge it is hard to explain why would migrants need a diet richer in protein (from animal prey) and presumably poorer in lipids (Bairlein 2002). It should be noted that muscle-score was not a significant predictor of trophic level, suggesting that body condition or

composition were not strongly correlated with diet. Furthermore, given that differences were present throughout winter, well before spring migratory preparation (fattening) occurs, this alternative explanation seems less likely.

Our findings have implications for models tackling the conditions for the evolution and maintenance of migratory systems. For example, in a theoretical modelling exercise, Taylor & Norris (2007), based on empirical evidence from previous studies (e.g. Adriaensen & Dhondt 1990, Pérez-Tris & Tellería 2002) assumed that migrants experience reduced competitive ability during winter. The results of the present study suggest that (longer-distance) migrants may not necessarily be competitively inferior. Northern robins were not only able to occupy the same habitats as their southern conspecifics despite arriving later to Iberia (Bueno 1998) but they were at a higher trophic level due to a greater consumption of invertebrates. Robins are known to prefer animal to plant foods (Berthold 1976), and so northern individuals may not be faring worse than southern counterparts.

Conclusions

Wintering robins in Iberia have been conclusively shown to clearly segregate between habitats by age, sex and size classes, which has been mostly attributed to the competitive exclusion of weaker individuals from preferred sites by conspecific dominants (Figuerola et al. 2001, Tellería et al. 2001, Catry et al. 2004). The evidence from our work shows that such segregation does not apply to migrants from different regions, suggesting no important asymmetries in the resource holding potential of birds originating from different populations. However, although there are no reported differences in diet between sex and age classes, there is clear evidence for dietary differences between northern and southern individuals. These are perhaps more readily explained

by the transportation to the wintering grounds of specializations originating in the breeding areas, but more research is needed to better evaluate the factors underlying these fascinating patterns, and to understand whether this particular type of ecological matching is a frequent trait in migratory species.

ACKNOWLEDGEMENTS

Many people provided feather samples, particularly L. Demongin and associated French ringers, J. Arizaga and ringers from Txingudi, V. Bulyuk, T. Valkenburg, P. Tenreiro, S. Marques, F. Sabino, R. Brito, A. C. Pereira, M. Araújo, F. Barros and H. Cardoso. Falsterbo Bird Station and its staff helped with sampling (carried out under license number M64-05). H. Alonso helped with fieldwork and Thomas Bodey in the lab. J. Robalo, V. Almada and S. Chenu provided help for molecular sexing. Two anonymous reviewers made useful comments on earlier versions of the manuscript. Fundação para a Ciência e a Tecnologia-Portugal provided grants SFRH/BD/30724/2006 and PEst-OE/MAR/UI0331/2011, as well as SFRH/BPD/40667/2007 (to JMN). Isotope ratio measurements were possible thanks to a Natural Environment Research Council LSMSF grant (EK106-01/07). To ICNF and Parque Natural da Arrábida for permission to carry out this research and to Companhia das Lezírias (particularly Eng. R. Alves) for allowing access to Charneca. All applicable institutional and/or national guidelines for the care and use of animals were followed.

LITERATURE CITED

Adriaensen F, Dhondt AA (1990) Population dynamics and partial migration of the European robin *Erithacus rubecula* in different habitats. *J Anim Ecol* 59:1077-1090.

411 Alves JA, Lourenço PM, Piersma P, Sutherland WJ, Gill JA (2010) Population overlap and
 412 habitat segregation in wintering black-tailed godwits *Limosa limosa*. Bird Study 57:381-
 413 391.

414 Bairlein F (comp.) (1995) Manual of field-methods. European-African songbird migration
 415 network. Wilhelmshaven, Germany.

416 Bairlein F (2002) How to get fat: nutritional mechanisms of seasonal fat accumulation in
 417 migratory songbirds. Naturwissenschaften 89:1–10

418 Bell CP (2005) Inter- and intrapopulation migration patterns: ideas, evidence and research
 419 priorities. In: Greenberg R, Marra PP (eds.). Birds of two worlds. The Ecology and
 420 Evolution of Migration. John Hopkins University Press, Baltimore.

421 Berthold P (1976) The control and significance of animal and vegetable nutrition in omnivorous
 422 songbirds. Ardea 64:140-154.

423 Block BA, Teo SLH, Walli A, et al. (2005) Electronic tagging and population structure of
 424 Atlantic bluefin tuna. Nature 434:1121-1127

425 Blumenthal JM, Abreu-Grobois FA, Austin TJ, et al. (2009) Turtle groups or turtle soup:
 426 dispersal patterns of hawksbill turtles in the Caribbean. Mol Ecol 18:4841-4853.

427 Bueno JM (1998) Migracion e invernada de pequeños turdinos en la Peninsula Ibérica V.
 428 Petirrojo (*Erithacus rubecula*). Ardeola 45:193-200.

429 Campos AR, Catry P, Tenreiro P, Neto JM, Pereira AC, Brito R, Cardoso H, Ramos JA, Bearhop
 430 S, Newton J (2011a) How do Iberian robins *Erithacus rubecula* respond to seasonal
 431 flooding by conspecific migrants? Bird Study 58:435–442.

432 Campos AR, Catry P, Ramos J, Robalo J (2011b) Competition among European robins *Erithacus*
 433 *rubecula* in the winter quarters: sex is the best predictor of priority of access to
 434 experimental food resources. *Ornis Fenn* 88:226-233.

435 Catry P, Campos A, Almada V, Cresswell W (2004) Winter segregation of migrant European
 436 robins *Erithacus rubecula* in relation to sex, age and size. *J Avian Biol* 35:204-209.

437 Catry P, Costa H, Elias G, Matias R (2010) Aves de Portugal. *Ornitologia do Território*
 438 Continental. Assírio & Alvim, Lisboa.

439 Catry P, Dias MP, Phillips RA, Granadeiro JP (2013) Carry-over effects from breeding modulate
 440 the annual cycle of a long-distance migrant: an experimental demonstration. *Ecology*
 441 94:1230-1235

442 Cramp S (Ed.) (1988) The Birds of the Western Palearctic, Vol V. Oxford University Press,
 443 Oxford.

444 Cuadrado M (1997) Why are migrant robins *Erithacus rubecula* territorial in winter?: the
 445 importance of the anti-predation behaviour. *Ethol Ecol Evol* 9:77-88.

446 de la Hera I, Pérez-Tris J, Tellería JL (2012) Habitat distribution of migratory and sedentary
 447 blackcaps *Sylvia atricapilla* wintering in southern Iberia: a morphological and
 448 biogeochemical approach *J Avian Biol* 43:333–340.

449 Debusse M, Isenmann P 1985. Frugivory of transient and wintering European robins *Erithacus*
 450 *rubecula* in a Mediterranean region and its relationship with ornithochory. *Holarctic Ecol*
 451 8:157-163.

452 Duijns S, van Dijk JGB, Spaans B, Jukema J, de Boer WF, Piersma T (2009) Foraging site
 453 selection of two subspecies of bar-tailed godwit *Limosa lapponica*: time minimizers
 454 accept greater predation risk than energy minimizers. *Ardea* 97:51-59.

455 Evans KL, Newton J, Gaston KJ, Sharp SP, McGowan A, Hatchwell BJ (2012) Colonisation of
 456 urban environments is associated with reduced migratory behaviour, facilitating
 457 divergence from ancestral populations. *Oikos* 121:634-640.

458 Figuerola J, Jovani R, Sol D (2001) Age-related habitat segregation by robins *Erithacus rubecula*
 459 during the winter. *Bird Study* 48:252-255.

460 Fonteneau F, Paillisson J-M, Marion L 2009 Relationships between bird morphology and prey
 461 selection in two sympatric great cormorant *Phalacrocorax carbo* subspecies during
 462 winter. *Ibis* 151:286-298

463 Fretwell SD, Lucas HL Jr. (1970) On territorial behavior and other factors influencing habitat
 464 distributions in birds.. *Acta Biotheoret* 19:16-36.

465 Fox AD, Christensen TK, Bearhop S, Newton J (2007) Using stable isotope analysis of multiple
 466 feather tracts to identify moulting provenance of vagrant birds: a case study of Baikal teal
 467 *Anas formosa* in Denmark. *Ibis* 142:622-625.

468 Gosler AG (1991) On the use of greater covert moult and pectoral muscle as measures of
 469 condition in passerines with data for the great tit *Parus major*. *Bird Study* 31:1-9.

470 Greenberg R (1986) Competition in migrant birds in the nonbreeding season. In: Johnston RF
 471 (ed) *Current Ornithology* 3. Plenum Press, New York and London, pp 281-307.

472 Greenberg R, Gonzalez CE, Bichier P, Reitsma R (2001) Nonbreeding habitat selection and
 473 foraging behavior of the black-throated green warbler complex in Southeastern Mexico.
 474 *Condor* 103:31-37.

475 Griffiths R, Double MC, Orr K, Dawson RJG (1998) A DNA test to sex most birds. *Mol Ecol*
 476 7:1071- 1075.

477 Gunnarsson TG, Gill JA, Newton J, Potts PM, Sutherland WJ (2005) Seasonal matching of
 478 habitat quality and fitness in a migratory bird. *Proc R Soc B* 272:2319-2323.

479 Herrera CM (1977) Ecología alimenticia del petirrojo (*Erithacus rubecula*) durante su invernada
 480 en encinares del sur de España. *Doñana, Acta Vert* 4:35-59.

481 Herrera CM (1978) Individual dietary differences associated with morphological variation in
 482 robins *Erithacus rubecula*. *Ibis* 120:542-545.

483 Herrera CM (1998) Long-term dynamics of Mediterranean frugivorous birds and fleshy fruits: a
 484 12-year study. *Ecol Monogr* 68:511-538.

485 Hjernquist MB, Veen T, Font L, Klaassen (2009) High individual repeatability and population
 486 differentiation in stable isotope ratios in winter-grown collared flycatcher *Ficedula*
 487 *albicollis* feathers. *J Avian Biol* 40:102-107.

488 Hobson KA and Bairlein F (2003) Isotopic fractionation and turnover in captive garden warblers
 489 (*Sylvia borin*): implications for delineating dietary and migratory associations in wild
 490 passerines. *Can J Zool* 81:1630–1635

491 Hobson KA (2008) Applying isotopic methods to tracking animal movements. In: Hobson KA,
 492 Wassenaar LI (eds) *Tracking Animal Migration with Stable Isotopes*. Academic Press,
 493 London.

494 Hobson KA, Wassenaar LI (2008) *Tracking Animal Migration with Stable Isotopes*. Academic
 495 Press, London.

496 Hobson KA, Bowen G, Wassenaar LI, Ferrand Y, Lormee H (2004) Using stable hydrogen
 497 isotope measurements of feathers to infer geographical origins of migrating European
 498 birds. *Oecologia* 141:477-488.

499 Inger R, Bearhop S (2008) Applications of stable isotope analyses to avian ecology. *Ibis*
 500 150:447-461.

501 Karlsson L, Persson K, Pettersson J, Walinder G (1988) Fat-weight relationships and migratory
 502 strategies in the robin *Erithacus rubecula* at two stop-over sites in South Sweden. *Ring*
 503 *Migr* 9:160-168.

504 Lockwood R., Swaddle JP, Rayner JMV (1998) Avian wing tip shape reconsidered: wingtip
 505 shape indices and morphological adaptations to migration. *J Avian Biol* 29:273-292.

506 Madigan D, Raftery AE (1994) Model selection and accounting for model uncertainty in
 507 graphical models using Occam's window. *J Am Stat Assoc* 89:1535–1546

508 Marion L (1995) Where two subspecies meet: origin, habitat choice and niche segregation of
 509 cormorant *Phalacrocorax c. carbo* and *P. c. sinensis* in the common wintering area
 510 (France), in relation to breeding isolation in Europe. *Ardea* 83:103–114.

511 Marra PP (2000) The role of behavioral dominance in structuring patterns of habitat occupancy
 512 in a migrant bird during the nonbreeding season. *Behav Ecol* 11:299-308.

513 McCormack JE, Smith TB (2008) Niche expansion leads to small-scale adaptive divergence
 514 along an elevation gradient in a medium-sized passerine bird. *Proc R Soc B* 275:2155-
 515 2164.

516 Newton I (2008) The migration ecology of birds. Academic Press, Amsterdam.

517 Paradis E, Baillie SR, Sutherland WJ, Gregory RD 1998. Patterns of natal and breeding dispersal
 518 in birds. *J Anim Ecol* 67:518-536.

519 Parnell AC, Inger R, Bearhop S & Jackson AL (2010) Source partitioning using stable isotopes:
 520 coping with too much variation. *PLoS ONE* 5:e9672.

521 Pérez-Tris J, Carbonell R, Tellería JL (2000) Identificación e importancia poblacional de los
 522 petirrojos *Erithacus rubecula* locales durante la invernada en el sur de España. *Ardeola*
 523 47:9-18.

524 Pérez-Tris J, Tellería JL (2002) Migratory and sedentary blackcaps in sympatric non-breeding
 525 grounds: implications for the evolution of avian migration. *J Anim Ecol* 71:211-224.

526 Peterson AT (1993) Adaptive geographical variation in bill shape of scrub jays (*Aphelocoma*
 527 *coerulescens*). *Am Nat* 142:508-527.

528 Peterson AT & Holt RD (2003) Niche differentiation in Mexican birds: using point occurrences
 529 to detect ecological innovation. *Ecol Let* 6:774-782.

530 Price T, Gross S (2005) Correlated evolution of ecological differences among the Old World leaf
 531 warblers in the breeding and nonbreeding seasons. In: Greenberg R, Marra PP (eds.)
 532 Birds of two worlds. The Ecology and Evolution of Migration. John Hopkins University
 533 Press, Baltimore.

534 R Development Core Team (2012) R: a language and environment for statistical computing.
 535 Vienna: R Foundation for Statistical Computing. <http://www.R-project.org/>

536 Raftery A, Hoeting J, Volinsky C, Painter I, Yeung KY (2013) BMA: Bayesian Model
 537 Averaging, R package version 3.16.2.3, <http://CRAN.R-project.org/package=BMA>

538 Rayner MJ, Hauber ME, Steeves TE, Lawrence HA, Thompson DR, Sagar PM, Bury SJ,
 539 Landers TJ, Phillips RA, Ranjard L, Shaffer SA (2011) Contemporary and historical
 540 separation of transequatorial migration between genetically distinct seabird populations.
 541 *Nature Com* 2: article number 332.

542 Ricklefs RE (1992) The megapopulation: a model of demographic coupling between migrant
 543 and resident landbird populations. In: Hagan III JM & Johnston DW (eds) *Ecology and*

544 Conservation of Neotropical Migrant Landbirds. Smithsonian Institution Press,
 545 Washington.

546 Svensson L (1992) Identification Guide to European Passerines. Svensson, Stockholm.

547 Taylor CM, Norris DR (2007) Predicting conditions for migration: effects of density dependence
 548 and habitat quality. *Biol Let* 3:280-283.

549 Tellería JL, Pérez-Tris J (2004) Consequences of the settlement of migrant European robins
 550 *Erithacus rubecula* in wintering habitats occupied by conspecific residents. *Ibis* 146:258-
 551 268.

552 Tellería JL, Blázquez M, de la Hera I, Pérez-Tris J (2013) Migratory and resident blackcaps
 553 *Sylvia atricapilla* wintering in southern Spain show no resource partitioning. *Ibis*
 554 155:750-761.

555 Tellería JL, Pérez-Tris J, Ramírez A, Fernández-Juridic E, Carbonell R (2001) Distribution of
 556 robins *Erithacus rubecula* in wintering grounds: effects of conspecific density, migratory
 557 status and age. *Ardea* 89:363-373.

558 Tobias J (1997) Asymmetric territorial contests in the European robin: the role of settlement
 559 costs. *Anim Behav* 54:9-21.

560 Wassenaar LI, Hobson KA (1998) Natal origins of migratory monarch butterflies at wintering
 561 colonies in Mexico: new isotopic evidence. *Proc Natl Acad Sci* 95:15436-15439.

562 Wassenaar LI, Hobson KA (2003) Comparative equilibration and online technique for
 563 determination of non-exchangeable hydrogen of keratins for use in animal migration
 564 studies. *Isotopes in Environmental and Health Studies* 39:211-217.

565 Wunder MB, Kester CL, Knopf FL, Rye RO (2005) A test of geographic assignment using
 566 isotope tracers in feathers of known origin. *Oecologia* 144:607-617.

567

568

569

570 Table 1. Effects of several predictors on $\delta^{15}\text{N}$ levels in blood of robins, using $\delta^2\text{H}$ (a) and C2 (b)
571 as proxy for geographic origin, estimated by Bayesian model averaging. Values represent
572 posterior probability of each variable has a non-zero coefficient in the model (PP), averaged
573 coefficients (conditional on the variable being in the model) and corresponding standard
574 deviation. Variables with high relevance are represented in bold.

575

| (a) Seventeen models averaged, the best 5 of which account for a cumulative probability of 0.67 | Posterior Probability (PP) | Averaged Coefficients | SD of averaged coefficients |
|--|----------------------------------|--------------------------|--------------------------------|
| Intercept | 100 | 2.115 | 1.049 |
| Site (habitat) | 100 | 1.532 | 0.209 |
| Age | 11.5 | -0.215 | 0.235 |
| Sex | 7.5 | -0.161 | 0.224 |
| $\delta^2\text{H}$ | 83.2 | -0.023 | 0.009 |
| Month (February) | 100 | -0.648 | 0.866 |
| Month (After February) | 100 | 0.654 | 1.043 |
| Muscle | 8.1 | -0.136 | 0.201 |
| Month*Site | 0 | 0.000 | 0.000 |
| Month*Age | 0 | 0.000 | 0.000 |
| Month*Sex | 0 | 0.000 | 0.000 |
| Site*Age | 5.7 | -0.088 | 0.333 |
| Site*Sex | 7.6 | -0.319 | 0.435 |

| | | | |
|---------|------|-------|-------|
| Age*Sex | 31.4 | 0.438 | 0.259 |
|---------|------|-------|-------|

576

| (b) Seven models averaged, the best 5 of which account for a cumulative probability of 0.67 | Posterior Probability | Averaged Coefficients | SD of averaged coefficients |
|--|--------------------------|--------------------------|--------------------------------|
| Intercept | 100 | 8.310 | 1.274 |
| Site (habitat) | 100 | 1.371 | 0.201 |
| Age | 7.1 | -0.106 | 0.201 |
| Sex | 7 | -0.101 | 0.201 |
| C2 | 100 | -2.495 | 0.690 |
| Month (February) | 100 | -0.854 | 0.712 |
| Month (After February) | 100 | 0.490 | 0.714 |
| Muscle | 6.6 | 0.074 | 0.208 |
| Month*Site | 0 | 0.000 | 0.000 |
| Month*Age | 0 | 0.000 | 0.000 |
| Month*Sex | 0 | 0.000 | 0.000 |
| Site*Age | 6.3 | -0.065 | 0.314 |
| Site*Sex | 6.3 | -0.050 | 0.349 |
| Age*Sex | 11.7 | 0.278 | 0.247 |


577

578


Fig. 1. Box-plot of $\delta^2\text{H}$ values of robins sampled during the breeding season in Portugal (N=69), France (N = 20), Scandinavia (N = 36) and Kaliningrad enclave, Russia (N = 7) and in winter in Iberia (n=170).

Fig 2. Relationship between (A) $\delta^2\text{H}$ in feathers (grown in spring/summer) or (B) wing-pointedness index (C_2), and $\delta^{15}\text{N}$ in whole blood of wintering robins. See Table 1 for statistics.

587 **Fig. 1**



590 **Fig. 2**



591

592

593

594

595